

KU KANSAS GEOLOGICAL SURVEY
The University of Kansas


MAP M-28 (Revised)

SURFICIAL GEOLOGY OF FINNEY COUNTY, KANSAS

Original geology by William C. Johnson and Alan F. Arbogast (1993)
Playa deposits and Arkansas River terrace deposits by William C. Johnson and Terri L. Woodburn

2010

Computer compilation and cartography by Jorgina A. Ross and Siew Phing Lee (2003)
Cartographic revisions by Christopher R. Becker, Nathaniel E. Haas, Scott T. Klopfenstein, R. Zane Price, and John W. Dunham (2010)



CENOZOIC ROCKS

Unconsolidated floodplain alluvium - Alluvium is found in the Arkansas River valley, Pawnee River valley, and valleys of other small streams in the county. Alluvial sediment is primarily coarse gravel, sand, and silt with a thickness that ranges from 30 to 50 feet in the Arkansas River valley, and from 12 to 30 feet in the Pawnee River valley.

Alluvial terrace deposits - Terrace deposits within the county occur along the Arkansas River, and the Pawnee River and its tributaries. Coarse gravel dominates the fill, with some sand and silt. The Ogallala Formation is the primary source. Age of the terrace fill ranges from the late Pleistocene to at least the middle Holocene. Three terrace levels are found in the county: Q₁ along the Pawnee River and along the main channel of the Arkansas River, and Q₂ and Q₃ along the Arkansas River valley in the western part of the county. The principal terrace of the Arkansas River (Q₁) lies 15 to 22 feet above the floodplain alluvium. Thickness of the other terrace deposits can reach 60 feet or more.

Upland intermittent lake (playa) deposits - Shallow basins, also known as playas, lagoons, or badland valleys, have developed in the upland basin deposits north and south of the Arkansas River valley. The origin of these features is usually attributed to wind deflation, animal activity, dissolution, or some combination of these processes. Age of the features ranges from at least the early Holocene to the historic period. The basins range in size from less than an acre to hundreds of acres. Fill within these basins has an average thickness of 7 feet and consists of reworked silt and fine sand from the loess. In the larger basins, a carbonaceous layer (caliche) typically develops a few feet below the basin floor.

Dune sand - Sand dunes occur immediately south of the Arkansas River and in small regions north of the Arkansas River and near the Pawnee River. The sand is derived from the Pleistocene terrace of the Arkansas River valley. Dunes reach a height of about 70 feet.

Sheet sand - Sand occurring in sheets, or subsoil underlying swales and swales, is expressed immediately south of the Arkansas River and in some of the basins where dune sand is found south of the river. The origin of the sand is the Pleistocene terrace of the Arkansas River valley. Sheet sand may reach a thickness of about 20 feet.

Active sand - Areas of blow sand as they existed at the time of mapping. Active sand areas are typically expressed as blowouts within the sand dune map units.

Loess - Wind-deposited silt, with minor amounts of clay and fine sand, comprises the loess, which marks the uplands in the county north of the Arkansas River valley and south of the sand region. The loess is calcareous and a buff color, and ranges in age from the late Pleistocene to the late Holocene. Loess thickness averages about 11 feet.

Ogallala Formation - Ogallala sediments are believed to be primarily Miocene in age and are composed of calcareous gravel, sand, silt, and clay deposited by streams transporting sediments eastward from the Rocky Mountains. Lenses of volcanic ash are found within the formation in this county. The Ogallala is commonly capped by thick calcareous beds known as the "mottled beds." These mottled beds, in turn, typically have a thin limestone bed capping them. The Ogallala mapped in Finney County also includes the early Pleistocene and older alluvial units of the same Rocky Mountain gravel and sand materials. Within the county, the Ogallala crops out along the north side of the Arkansas River and south side of the Pawnee River and along the higher bluffs in the northeastern part of the county. Most of the county is underlain by the Ogallala, where it reaches a thickness of greater than 500 feet (with thin areas). In the northeast part, surface exposures occur with a thickness of up to 47 feet.

Fort Hays Limestone Member - Cretaceous Fort Hays Limestone, a member of the Noboria Chalk, is composed of thick beds of chalk separated by thin beds of cherty shale. Exposures are found along the tributary streams of the Pawnee River in the northeastern part of the county, with an outcrop thickness of up to 52 feet.

Blue Hill Shale Member - Cretaceous Blue Hill Shale is the oldest rock cropping out in the county. It crops out along the Pawnee River and in tributaries in the northeastern part of the county. This member is a bluish-black, noncalcareous clay shale with an outcrop thickness of up to 92 feet.

Geologic Unit Boundaries

- Observed contact
- Inferred contact

Hydrology and Topography

- Intermittent stream
- Perennial stream
- Stream
- Intermittent water body
- Perennial water body
- Stream
- Flowline contour (Observed contour)
- Elevation contour (Observed contour)
- Depression contour (Observed contour)
- Flowline contour (Inferred contour)
- Elevation contour (Inferred contour)
- Depression contour (Inferred contour)
- Levee

Resource Development

- Oil
- Gas
- Water
- Open-pit gravel pit
- Open-pit sand pit

Index to 1:24,000-scale USGS quadrangle maps

This index shows the names and locations of the 36 USGS 7.5-minute 1:24,000-scale quadrangles used in the digital compilation of the Finney County map. The geology was mapped on the grid using these quadrangle maps.

The 2010 revision consists of two additions: (1) terrace systems along the Arkansas River that had previously been mapped as alluvium and (2) playa deposits that were not shown on the original geologic map.

Elevation contours are presented for general reference. They are taken from the USGS Digital Line Graph (DLG) files compiled from base maps at a scale of 1:50,000. In some areas the contours from the DLG may not show potential local variations (such as the complex of gullies, swales, and pits). Missing portions on the map will typically indicate topographic variations more accurately than the generalized contour lines. Reported elevations of an existing water contour line should be interpreted as an indication that the mapped rock unit is maintaining a relatively constant elevation above a generalized contour.

The geology was mapped using the ArcGIS system developed by ERI Environmental Systems Research Institute, Inc. (2003).

Peak and highest elevations shown on the map are summarized by data from the Kansas Department of Transportation (KDOT) and other sources. U.S. Department of Agriculture - Farm Service Agency (FSA) National Agriculture Imagery Program (NAIP) imagery also was used to check modifications.

Shaded relief is based on a 1:50,000-scale digital elevation model (DEM) with 15-m resolution. The shaded relief was generated using a 3D DEM. The DEM was processed using a 3D DEM processing algorithm to generate shaded relief. The shaded relief was then converted to a hillshade, a multidirectional shaded-relief image using angles of observation from 0°, 22.5°, 45°, and 67.5° azimuths, each 45° above the horizon, with a 4x vertical exaggeration.

This map was produced using the ArcGIS system developed by ERI Environmental Systems Research Institute, Inc. (2003).

The Kansas Geological Survey does not guarantee this map to be free from errors or omissions and does not assume any responsibility for interpretations made from the map or associated hard copies.

Generalized Geology of Kansas

- Quaternary System
- Neogene System
- Cambrian-Ordovician System
- Paleozoic System
- Permian System
- Carboniferous System
- Mississippian Subsystem

Geologic Unit Boundaries

- Observed contact
- Inferred contact

Hydrology and Topography

- Intermittent stream
- Perennial stream
- Stream
- Intermittent water body
- Perennial water body
- Stream
- Flowline contour (Observed contour)
- Elevation contour (Observed contour)
- Depression contour (Observed contour)
- Flowline contour (Inferred contour)
- Elevation contour (Inferred contour)
- Depression contour (Inferred contour)
- Levee

Resource Development

- Oil
- Gas
- Water
- Open-pit gravel pit
- Open-pit sand pit

Location Diagram

SCALE 1:50,000

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100

1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17 18 19 20 21 22 23 24 25 26 27 28 29 30 31 32 33 34 35 36 37 38 39 40 41 42 43 44 45 46 47 48 49 50 51 52 53 54 55 56 57 58 59 60 61 62 63 64 65 66 67 68 69 70 71 72 73 74 75 76 77 78 79 80 81 82 83 84 85 86 87 88 89 90 91 92 93 94 95 96 97 98 99 100